

Technical documentation

Product name:	qRE Echinacea angustifolia DC., roots
Substance:	Echinacea angustifolia DC., roots dry extract
Plant source common names:	en: Narrow-leaved purple coneflower; fr: Echinacée angustifolia
Reference:	E0112
Packaging:	100 mg in a 1.5 ml borosilicate amber vial
Storage conditions:	Keep container closed. Protect from light and moisture. Keep inferior to -15 °C.
Retest:	12 months

Botanical identification of plant source

Plants in our botanical garden are identified and a herbal voucher is prepared by an expert botanist.
Each batch collected for extraction is verified and identified.

Reference: Flora of North America http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=242416463

Method of production of dry extract

Whole plant or plant parts are collected, freeze-dried and coarsely ground. Extraction is performed by maceration in 50 % (v/v) aqueous ethanol (v/v) for 48 hours at room temperature. Ethanol is then evaporated under reduced pressure at less than 40 °C and the aqueous residue is freeze-dried.

Residual water content measurement is done by Karl Fischer titration.

Organoleptic characteristics of dry extract

Colour: Emerald green

Odour: Non characteristic

Form: Fine powder

Recommended methods for use

Weight a precise weight of qRE and solubilise in the recommended solvent at the concentration indicated in the HPLC or HPTLC method described in this document.

Sonicate for 90 seconds (70 W).

Filter on a 0.45 µm PVDF membrane and put the resulting solution into HPLC dispenser or apply on the HPTLC plate.

Dose and analyse your extract with qRExtract using the HPLC / HPTLC methods described in this document or using your own methods.

HPTLC

Detection of chicoric acid and echinacoside

Layer: 10 × 10 cm HPTLC Nano-Sil-20 UV 254 (Carl Roth ref. N084.1)

Thin layer conditionnement: 1 h at room temperature and 33 % relative humidity


Elution solvent:	<u>Elution solvent compound</u>	<u>Volume (ml)</u>
	ethyl acetate	100
	water	27
	formic acid	11
	acetic acid	11

Developing distance: 70 mm from the lower edge

Initial spot volume and concentration:

chicoric acid:	1 µl of a 0.2 % (w/v) solution in ethanol 96 %
qRE:	5 µl of a 1.5 % (w/v) solution in 50 % (v/v) aqueous ethanol
echinacoside:	1 µl of a 0.02 % (w/v) solution in ethanol 96 %




Reagent mixture: Natural products - polyethylene glycol reagent (NP/PEG)
Preparation: Dissolve 0.25 g of diphenylboric acid 2-aminoethylester and 1.25 g of polyethylene glycol 400 in 25 mL of methanol.
Dip the plate in the reagent mixture and dry for 15 minutes at room temperature.
Expose to UV light at 365 nm.



HPLC

Precolumn:	Ascentis® Express C18 0.5 cm × 3.0 mm 2.7 µm		
Column:	Ascentis® Express C18 15 cm × 3.0 mm 2.7 µm		
Sample:	8 µl 2.17 % qRE® (w/v) solution in 50 % (v/v) aqueous ethanol		
Flow:	0.45 ml/min		
Temperature:	25 °C		
Mobile phase:	A: 0.1 % formic acid (v/v) in water B: 0.1 % formic acid (v/v) in acetonitrile		
Detection:	Diode Array Detector, 210 nm		
Gradient:	Time (mn)	A %	B %
	0	97	3
	40	76	24
	50	52	48
	80	15	85

Quantified substances

Compound	CAS No	2D Structure	Peak No
Echinacoside	82854-37-3		2
Undeca-2,4-diene-8,10-diynoic acid isobutylamide	13891-74-2		3
Pentadeca-2,9-diene-12,14-diynoic acid isobutylamide	NA		4
Unknown	NA	NA	1, 5